

## Greybody Factors in the $AdS_3$ /CFT Correspondence

H.W. Lee and Y. S. Myung

*Department of Physics, Inje University, Kimhae 621-749, Korea*

### Abstract

We propose a new method to calculate the greybody factor in the  $AdS_3$ . This is based on both the non-normalizable modes of a test field( $\Phi_i$ ) and  $AdS_3$ /CFT correspondence. Such non-normalizable modes serve as classical, non-fluctuating background and encode the choice of operator insertion( $\mathcal{O}^i$ ) in the boundary. Actually specifying the boundary condition at infinity of  $AdS_3$  corresponds to turning on these non-normalizable modes. Hence we can calculate the greybody factor of test fields without the Dirichlet(or Neumann) boundary condition. The result is consistent with those of the boundary CFT and effective string calculations.

Recently the *AdS/CFT* correspondence has attracted much interest [1–3]. This is based on the duality relation between the string theory on  $AdS_{d+1} \times M$  and a conformal field theory living on the  $d$ -dimensional boundary( $\mathcal{B}$ ) of the *AdS* sector. The relevant relation between string theory in the bulk and field theory on the boundary is

$$\mathcal{Z}_{\text{eff}}(\Phi_i) = e^{iS_{\text{eff}}(\Phi_i)} = \langle T e^{i \int_{\mathcal{B}} \Phi_{b,i} \mathcal{O}^i} \rangle, \quad (1)$$

where  $S_{\text{eff}}$  is the effective action in the bulk,  $\Phi_{b,i}$  is the test field  $\Phi_i$  on the boundary, and  $T$  is the time-ordered product in the boundary. There is a one-to-one correspondence between the local operator( $\mathcal{O}^i$ ) of the boundary CFT and the bulk(test) field( $\Phi_i$ ) in  $AdS_3$ . For example, the test fields $\{\Phi_i\}$  are those in the D1-D5 brane system: a free scalar( $\phi$ ) which couples to (1,1) operator in the boundary; two fixed scalars ( $\nu, \lambda$ ) to (2,2), (3,1), and (1,3) operators [4,5]; two intermediate scalars ( $\eta, \xi$ ) to (1,2) and (2,1) operators [6]. The expectation value  $\langle \dots \rangle$  is taken in the CFT with  $\Phi_{b,i}$  as a source. From the boundary CFT, we can derive the bulk equation of  $\Phi_i$  with mass( $m$ ) and spin( $s$ ) [7,8]. Here we take  $AdS_3 \times S^3$  as a relevant model. This is so because the (D-brane) 5D black hole becomes  $AdS_3 \times S^3$  near horizon. Using the *AdS*<sub>3</sub>/*CFT* correspondence, one can obtain much information of the dynamical aspects(greybody factor = absorption cross section) of this black hole. Actually the  $AdS_3 \times S^3$  is an exact solution of string theory and there is an exact CFT on its boundary at spatial infinity [9].

Gubser has derived the general formular for the greybody factor by using the effective string model [9]:

$$\begin{aligned} \sigma_{\text{abs}} = & \frac{2(h_R + h_L - 1)^2 (2\pi l T_L)^{2h_L-1} (2\pi l T_R)^{2h_R-1}}{\omega \Gamma(2h_L) \Gamma(2h_R)} \\ & \times \sinh\left(\frac{\omega}{2T_H}\right) \left| \Gamma(h_L + i\frac{\omega}{2\pi T_L}) \Gamma(h_R + i\frac{\omega}{2\pi T_R}) \right|^2. \end{aligned} \quad (2)$$

Also Teo recovered the same form by using the boundary CFT [10]. In the boundary CFT calculation, the Poincaré coordinates ( $ds^2/l^2 = (-d\tilde{t}^2 + d\tilde{x}^2 + dz^2)/z^2 + d\Omega_3^2$ ) are used for studying the region from the  $z = l$  to the near horizon ( $z = \infty$ ). The boundary was taken to be at  $z = l$  rather than infinity( $z = 0$ ). He argued that this choice bypasses the problem

of field, which becomes divergent at infinity of  $AdS_3$ . The key information was encoded in the two-point function for  $\mathcal{O}^i$ ,

$$\langle \mathcal{O}(\mathbf{X}) \mathcal{O}(\mathbf{Y}) \rangle = \frac{2(h_L + h_R - 1)2l^{2(h_L + h_R) - 3}}{\pi |\mathbf{X} - \mathbf{Y}|^{2(h_L + h_R)}}. \quad (3)$$

In this paper, we calculate the greybody factor of a set of test fields by using their non-normalizable mode and  $AdS_3$ /CFT correspondence. The normalizable and non-normalizable modes emerge naturally either from a direct solution of the wave equation (5) or from the field representation of the  $AdS_3$  isometry group  $(SL(2, R)_L \times SL(2, R)_R)$  [11]. The former propagates in the bulk and corresponds to physical state, whereas the latter plays a role of classical, non-fluctuating background. We stress that the non-normalizable mode encodes the choice of operator insertion in the boundary. In the conventional study of  $AdS_3$ , the boundary condition is crucial for obtaining the sensible result [12]. This is so because the  $AdS_3$  is not globally hyperbolic, so that information may enter or exit from the boundary at infinity. There are the Dirichlet or Neumann boundary condition at infinity. If one requires the Dirichlet condition, the cross section can be calculated only for  $m^2 l^2 < -3/4$  [13]. Further the stability on the  $AdS_3$  requires  $m^2 l^2 \geq -1$  [3]. Hence if one takes the stability and the Dirichlet condition seriously, the conventional method can be applied for the test field belonging to  $-1 \leq m^2 l^2 < -3/4$ . Here includes the tachyon with  $m^2 l^2 = -1$  [7], which couples to minimal weight primary operator  $(1/2, 1/2)$ . However, the relevant fields  $\{\Phi_i\}$  are a free scalar field  $(\phi)$  with  $m^2 l^2 = 0$ , fixed scalars  $(\nu, \lambda)$  with  $m^2 l^2 = 8$ , and intermediate scalars  $(\eta, \xi)$  with  $m^2 l^2 = 3$ . Those remain unsolved in the conventional approach. Hence we have to develop a new calculation scheme to cover the relevant fields. This is just our method based on the non-normalizable modes [14–16]. In this case, specifying the boundary condition at spatial infinity corresponds to introducing the non-normalizable modes [11]. That is, the non-normalizable solution to the equation of motion corresponds to an operator insertion at infinity and a particular choice of boundary conditions. Hence, using this method, one can calculate the greybody factors of relevant fields without the additional Dirichlet or Neumann boundary condition.

We start with the effective action for a test field  $\Phi$  with mass  $m$  [3,11]

$$S_{\text{eff}} = \frac{1}{2} \int_{AdS_3} d^3x \sqrt{-g} \left[ (\nabla\Phi)^2 + m^2\Phi^2 \right], \quad (4)$$

where  $m^2 = (h_L + h_R)(h_L + h_R - 2)/l^2$  according to the  $AdS_3$ /CFT correspondence [3,8,11].

The equation of motion leads to

$$\nabla^2\Phi - m^2\Phi^2 = 0. \quad (5)$$

The BTZ black hole background (locally,  $AdS_3$ ) is given by [17]

$$\bar{g}_{\mu\nu} = \begin{pmatrix} -(M - r^2/l^2) & -J/2 & 0 \\ -J/2 & r^2 & 0 \\ 0 & 0 & f^{-2} \end{pmatrix} \quad (6)$$

with  $f^2 = r^2/l^2 - M + J^2/4r^2$ . The metric  $\bar{g}_{\mu\nu}$  is singular at  $r = r_{\pm}$ ,

$$r_{\pm}^2 = \frac{Ml^2}{2} \left\{ 1 \pm \left[ 1 - \left( \frac{J}{Ml} \right)^2 \right]^{1/2} \right\} \quad (7)$$

with  $M = (r_+^2 + r_-^2)/l^2$ ,  $J = 2r_+r_-/l$ . Here  $l \gg r_+ > r_-$ . For convenience, we list the Hawking temperature  $T_H$ , the area of horizon  $\mathcal{A}_H$ , and the angular velocity at the horizon  $\Omega_H$  as

$$T_H = (r_+^2 - r_-^2)/2\pi l^2 r_+, \quad \mathcal{A}_H = 2\pi r_+, \quad \Omega_H = J/2r_+^2. \quad (8)$$

Considering the  $t$  and  $x$ -translational symmetries of the background metric (6), one chooses the perturbation for a test field as

$$\Phi(t, \phi, r) = e^{-i\omega t} e^{i\mu\phi} \tilde{\Phi}(r). \quad (9)$$

Hence (5) becomes the equation [14]

$$\left[ f^2 \partial_r^2 + \left\{ \frac{1}{r} (\partial_r r f^2) \right\} \partial_r - \frac{J\mu\omega}{r^2 f^2} + \frac{\omega^2}{f^2} + \frac{M - \frac{r^2}{l^2}}{r^2 f^2} \mu^2 \right] \tilde{\Phi} - m^2 \tilde{\Phi} = 0, \quad (10)$$

Now we calculate the absorption cross section of the test field to obtain its dynamical information from the  $AdS_3$ /CFT correspondence. Since it is hard to find a solution to (10)

directly, we use a matching procedure. The spacetime is divided into two regions: the near region ( $r \sim r_+ \ll l$ ) and far region ( $r \gg l$ ). We now study each region in turn. For the far region ( $r \gg l$ ), the equation (10) becomes

$$\tilde{\Phi}_{\text{far}}'' + \frac{3}{r}\tilde{\Phi}_{\text{far}}' - \frac{\Lambda}{r^2}\tilde{\Phi}_{\text{far}} = 0. \quad (11)$$

Here  $\Lambda = m^2 l^2 + \epsilon$  is introduced with the small parameter  $\epsilon$  for the technical reason. First we find the far region solution

$$\tilde{\Phi}_{\text{far}}(x) = \frac{1}{x} \left( \alpha x^{\sqrt{1+\Lambda}} + \beta x^{-\sqrt{1+\Lambda}} \right) = \tilde{\Phi}_{\text{far}}^{\text{non-nor}} + \tilde{\Phi}_{\text{far}}^{\text{nor}} \quad (12)$$

with two unknown coefficients  $\alpha, \beta$  and  $x = r/l$ . We note that for the case of  $m^2 l^2 \geq 0$ , the first(second) term diverges(converges) at  $x = \infty$ , and thus turn out to be the non-normalizable(normalizable) modes [11]. We then calculate the ingoing flux at infinity and this is given by

$$\mathcal{F}_{\text{in}}(\infty) = -2\pi\sqrt{1+\Lambda}|\alpha - i\beta|^2. \quad (13)$$

Here we need both non-normalizable and normalizable modes to obtain the non-zero incoming flux at infinity. In order to obtain the near region behavior, we introduce the variable

$z = \frac{r^2 - r_+^2}{r^2 - r_-^2} = \frac{x^2 - x_+^2}{x^2 - x_-^2}$ ,  $0 \leq z \leq 1$ . Then Eq.(10) leads to

$$z(1-z)\frac{d^2\tilde{\Phi}}{dz^2} + (1-z)\frac{d\tilde{\Phi}}{dz} + \left( \frac{A_1}{z} - \frac{\Lambda/4}{1-z} + B_1 \right) \tilde{\Phi} = 0, \quad (14)$$

where  $A_1 = \left( \frac{\omega - \mu\Omega_H}{4\pi T_H} \right)^2$ ,  $B_1 = -\frac{r_-^2}{r_+^2} \left( \frac{\omega - \mu\Omega_H r_+^2/r_-^2}{4\pi T_H} \right)^2$ . The solution for Eq.(14) is given by the hypergeometric functions

$$\begin{aligned} \tilde{\Phi}_{\text{near}}(z) = & C_1 z^{-i\sqrt{A_1}} (1-z)^{(1-\sqrt{1+\Lambda})/2} F(a, b, c; z) \\ & + C_2 z^{i\sqrt{A_1}} (1-z)^{(1-\sqrt{1+\Lambda})/2} F(b-c+1, a-c+1, 2-c; z), \end{aligned} \quad (15)$$

where

$$\begin{aligned} a &= \sqrt{B_1} - i\sqrt{A_1} + (1 - \sqrt{1+\Lambda})/2, \\ b &= -\sqrt{B_1} - i\sqrt{A_1} + (1 - \sqrt{1+\Lambda})/2, \\ c &= 1 - 2i\sqrt{A_1}. \end{aligned}$$

and  $C_1$  and  $C_2$  are to-be-determined constants. At the near horizon( $r \sim r_+, z \sim 0$ ), (15) becomes

$$\begin{aligned}\tilde{\Phi}_{\text{near}}(0) &\simeq C_1 z^{-i\sqrt{A_1}} + C_2 z^{i\sqrt{A_1}} \\ &= C_1 \left( \frac{2x_+}{x_+^2 - x_-^2} \right)^{-i\sqrt{A_1}} e^{-i\sqrt{A_1} \ln(x-x_+)} + C_2 \left( \frac{2x_+}{x_+^2 - x_-^2} \right)^{i\sqrt{A_1}} e^{i\sqrt{A_1} \ln(x-x_+)}. \quad (16)\end{aligned}$$

Taking an ingoing mode at horizon, we have  $C_2 = 0$ . Hence the near region solution is

$$\tilde{\Phi}_{\text{near}}(z) = C_1 z^{-i\sqrt{A_1}} (1-z)^{(1-\sqrt{1+\Lambda})/2} F(a, b, c; z). \quad (17)$$

Now we need to match the far region solution (12) to the large  $r(z \rightarrow 1)$  limit of near region solution (17) in the overlapping region. The  $z \rightarrow 1$  behavior of (17) follows from the  $z \rightarrow 1-z$  transformation rule for hypergeometric functions. Using  $1-z \sim (x_+^2 - x_-^2)/x^2$  for  $r \rightarrow \infty$ , this takes the form

$$\tilde{\Phi}_{n \rightarrow f}(r) \simeq C_1 E_1 \frac{x^{\sqrt{1+\Lambda}}}{x} + C_1 E_2 \frac{x^{-\sqrt{1+\Lambda}}}{x}, \quad (18)$$

where

$$E_1 = \frac{\Gamma(1-2i\sqrt{A_1})\Gamma(\sqrt{1+\Lambda})(x_+^2 - x_-^2)^{(1-\sqrt{1+\Lambda})/2}}{\Gamma(\frac{1+\sqrt{1+\Lambda}}{2} + \sqrt{B_1} - i\sqrt{A_1})\Gamma(\frac{1+\sqrt{1+\Lambda}}{2} - \sqrt{B_1} - i\sqrt{A_1})}, \quad (19)$$

$$E_2 = \frac{\Gamma(1-2i\sqrt{A_1})\Gamma(-\sqrt{1+\Lambda})(x_+^2 - x_-^2)^{(1+\sqrt{1+\Lambda})/2}}{\Gamma(\frac{1-\sqrt{1+\Lambda}}{2} + \sqrt{B_1} - i\sqrt{A_1})\Gamma(\frac{1-\sqrt{1+\Lambda}}{2} - \sqrt{B_1} - i\sqrt{A_1})}. \quad (20)$$

Matching (12) with (18) leads to  $\alpha = C_1 E_1$  and  $\beta = C_1 E_2$ . Considering  $x_+^2 - x_-^2 \ll 1$  and  $m^2 l^2 \geq 0$ , then one finds  $\beta \ll \alpha$ . Thus the ingoing flux across the horizon is given by  $\mathcal{F}_{\text{in}}(0) = -8\pi\sqrt{A_1}(x_+^2 - x_-^2)|C_1|^2$ . This amounts to taking the flux of the non-normalizable mode effectively. At this stage, we wish to comment on the parameter  $\Lambda$ . Initially we introduce  $\Lambda = m^2 l^2 + \epsilon$  with the small parameter  $\epsilon$  in Eq.(11). This is so because  $E_2$  in (20) has a pole for integral  $\Lambda$  through  $\Gamma(-\sqrt{1+\Lambda})$ . Hence it is convenient to keep  $\Lambda$  near an integer value during the calculation and make it integer at the end. Hence for  $\mu = 0, \epsilon = 0 (\Lambda = m^2 l^2)$ , we can obtain the absorption coefficient

$$\begin{aligned}\mathcal{A} &= \frac{\mathcal{F}_{\text{in}}(0)}{\mathcal{F}_{\text{in}}(\infty)} = \frac{4\sqrt{A_1}(x_+^2 - x_-^2)}{\sqrt{1+m^2 l^2}} \frac{1}{|E_1|^2} \\ &= \frac{\omega \mathcal{A}_H}{\pi} \frac{(x_+^2 - x_-^2)^{(\sqrt{1+m^2 l^2}-1)}}{\Gamma(1+\sqrt{1+m^2 l^2})\Gamma(\sqrt{1+m^2 l^2})} \left| \frac{\Gamma(\frac{1+\sqrt{1+m^2 l^2}}{2} - i\frac{\omega}{4\pi T_L})\Gamma(\frac{1+\sqrt{1+m^2 l^2}}{2} - i\frac{\omega}{4\pi T_R})}{\Gamma(1-i\frac{\omega}{2\pi T_H})} \right|^2, \quad (21)\end{aligned}$$

where left and right temperatures are defined by

$$\frac{1}{T_{L/R}} = \frac{1}{T_H} \left( 1 \pm \frac{r_-}{r_+} \right). \quad (22)$$

The absorption cross-section is given by  $\sigma_{\text{abs}} = \mathcal{A}/\omega$  in three dimensions. For  $h_L = h_R$ , this takes the form

$$\begin{aligned} \sigma_{\text{abs}}^{AdS} = & \frac{2(h_L + h_R - 1)(2\pi l T_L)^{2h_L-1}(2\pi l T_R)^{2h_R-1}}{\pi\omega\Gamma(2h_L)\Gamma(2h_R)} \sinh\left(\frac{\omega}{2T_H}\right) \\ & \times \left| \Gamma\left(h_L - i\frac{\omega}{4\pi T_L}\right) \Gamma\left(h_R - i\frac{\omega}{4\pi T_R}\right) \right|^2. \end{aligned} \quad (23)$$

This is our key result. In the  $m^2 l^2 \rightarrow 0$  limit, (23) recovers the decay rate for the free scalar( $\phi$ ) which couples to (1,1) operator [15,18]

$$\Gamma_{\text{min}} = \frac{\sigma_{\text{abs}}^{\text{min}}}{e^{\frac{\omega}{T_H}} - 1} = \frac{\pi l^2 \omega}{(e^{\frac{\omega}{2T_R}} - 1)(e^{\frac{\omega}{2T_L}} - 1)}. \quad (24)$$

It is pointed out that the dilaton as a fixed scalar( $\nu$ ) which couples to (2,2) is physically propagating field in the BTZ back ground. In the limit of  $m^2 l^2 \rightarrow 8$ , our result (23) is that for the dilaton as was shown in Refs. [14,16]. For  $m^2 l^2 = 3$ , one can obtain the cross section for a new scalar which couples to the primary operator (3/2,3/2) [7].

In this  $AdS_3$ -calculation, there exists an ambiguity in determining normalization factor of the cross section for test field with  $m^2 l^2 > 0$  in comparison with the previous methods. In the case of semiclassical calculation with asymptotically flat space, there is no ambiguity in calculating the cross section [4,5]. This includes both  $M_5 \times S^1 \times T^4$  (5D black hole) and  $AdS_3 \times S^3 \times T^4$  near horizon with the asymptotically flat space [15,16]. Also, in the effective string approach one can calculate the cross section without arbitrariness [4,9]. However, as in the effective CFT method by Maldacena and Strominger [19], it seems that there is no known way to fix the normalization factor without appealing to string theory. To recover the effective string and boundary CFT results (2) correctly, we need a conversion factor ( $\pi(h_L + h_R - 1)$ ). As a definite example, we list the cross section of the fixed scalar( $\nu$ ) which couples to (2,2) operator. This is  $\sigma_{\text{abs}}^{\omega \rightarrow 0} = C \mathcal{A}_H(r_0/l)^4$ . Here  $C = 1/4$  for the semiclassical

calculation, effective string method, boundary CFT-calculation in (2), whereas  $C = 1/12$  for our  $AdS_3$ -calculation. However, there is no ambiguity for a free scalar with  $m^2 l^2 = 0$ .

Now we are in a position to discuss other fields which couple to chiral operators as (3,1), (1,3), (2,1), (1,2), (2,0), (0,2), (1,0), and (0,1). We remind the reader that the fixed scalar( $\lambda$ ) couples to (3,1) and (1,3) as well as (2,2). This has a discrepancy in the cross section between the semiclassical and effective string calculations [4,5]. Two intermediate scalars ( $\eta, \xi$ ) couple to (1,2) and (2,1) [6]. To obtain the general form of their greybody factors by using the semiclassical approach remains as open question still. Also gauge bosons couple to (2,0) and (0,2) and tachyons to (1,0) and (0,1) [7]. Those all carry non-zero spin( $s = h_L - h_R$ ) with the mass( $m^2 l^2 = (h_L + h_R)(h_L + h_R - 2)$ ). In order to study the wave equation including the non-zero spin, we may extract its information from the boundary CFT. We write down its equation on the boundary at the infinity as [8]

$$\left[ \nabla^2 + \frac{s^2}{l^2 \sinh^2 \rho} - m^2 \right] \Phi_s = 0 \quad (25)$$

in the global coordinates( $\tau, \phi, \rho$ ):

$$\frac{ds_g^2}{l^2} = -\cosh^2 \rho d\tau^2 + \sinh^2 \rho d\phi^2 + d\rho^2. \quad (26)$$

Its solution with  $\nu^2 = 1 + m^2 l^2$  is given by

$$\Phi_s = e^{-i(h_L + h_R)\tau - is\phi} \left[ \frac{\alpha}{(\cosh \rho)^{(1-\nu)/2}} + \frac{\beta}{(\cosh \rho)^{(1+\nu)/2}} \right]. \quad (27)$$

We note here that the first (the second) terms correspond to the non-normalizable (normalizable) modes. For large  $\rho$  ( $r \rightarrow \infty$ ),  $\Phi_s$  vanishes. Near horizon it takes a plane wave.

Introducing the Poincaré coordinates( $\omega^+ = \tilde{t} + \tilde{x}, \omega^- = \tilde{t} - \tilde{x}, z$ ) to rewrite (25) leads to

$$\left[ z^2 \left( 4\partial_+ \partial_- + \partial_z^2 - \frac{1}{z} \partial_z \right) - \frac{4s^2 z^2}{z^4 - 2(1 - \omega^+ \omega^-)z^2 + (1 + \omega^{+2})(1 + \omega^{-2})} - m^2 l^2 \right] \Phi_s = 0. \quad (28)$$

If  $s = 0$  and  $\Phi_{s=0} = e^{-i\omega\tilde{t} + i\mu\tilde{x}} z \tilde{\Phi}(\eta)$  with  $\eta = \sqrt{\omega^2 - p^2} z$ , the above reduces to the Bessel equation



$$\eta^2 \ddot{\Phi} + \eta \dot{\Phi} + [\eta^2 - \nu^2] \Phi = 0. \quad (29)$$

The solution for an integer  $\nu = h_L + h_R - 1$  is given by [7]

$$\Phi = e^{-i\omega\tilde{t} + i\mu\tilde{x}} \left[ \beta z J_\nu(\sqrt{\omega^2 - \mu^2} z) + \alpha z Y_\nu(\sqrt{\omega^2 - \mu^2} z) \right]. \quad (30)$$

The first term satisfies the Dirichlet boundary condition at  $z = 0$  ( $r \rightarrow \infty$ ) and corresponds to the normalizable mode. This may be used for the perturbative calculation. Although the latter does not satisfy the Dirichlet boundary condition, it corresponds the non-normalizable mode to specify another boundary condition. And thus this mode can be used to derive the greybody factor. However, turning on the spin-dependent term, we have a difficulty in solving Eq.(28). This is mainly due to being unable to separate those variables.

Finally, we wish to rewrite Eq.(25) in terms of the BTZ coordinates( $t, \phi, r$ ). For this purpose, we introduce the proper radial coordinates( $\tau, \phi, \rho$ ) [20,21]:

$$ds_P^2 = -\sinh^2 \rho (-r_+ d\tau + r_- d\phi)^2 + l^2 d\rho^2 + \cosh^2 \rho (-r_- d\tau + r_+ d\phi)^2 \quad (31)$$

with  $r^2 = r_+^2 \cosh^2 \rho - r_-^2 \sinh^2 \rho$  and  $\tau = t/l$  for the exterior of the BTZ black hole. Here we define  $\gamma$  and  $\delta$  as  $t$  and  $\phi$ ,

$$\gamma = \frac{-r_- \tau + r_+ \phi}{l}, \delta = \frac{-r_+ \tau + r_- \phi}{l}. \quad (32)$$

Then Eq.(25) can be rewritten as

$$\left[ l^2 \nabla_{\text{BTZ}}^2 - \frac{2s^2(r_+^2 - r_-^2)}{(r^2 - r_-^2) \cosh 2\gamma + (r^2 - r_+^2) \cosh 2\delta + (r_+^2 - r_-^2)} - m^2 l^2 \right] \Phi_s = 0. \quad (33)$$

Note that the spin term depends on  $(\phi, \tau)$  and takes the form of  $r^{-4}$ . This is very similar to Eq.(28) which tell us that its spin term depends on  $(\omega^+, \omega^-)$  and takes the form of  $z^{-4}$ . Near horizon( $r \rightarrow r_+$ ), Eq.(33) takes the following form with  $x, \tau$ , and  $j = J/l$ :

$$\left[ f^2 \partial_x^2 + \left\{ \frac{1}{x} \partial_x (x f^2) \right\} \partial - \frac{j}{x^2 f^2} \partial_\phi \partial_\tau - \frac{1}{f^2} \partial_\tau^2 - \frac{1}{f^2 x^2} (M - x^2) \partial_\phi^2 - \frac{s^2}{\cosh^2(x_+ \phi - x_- \tau)} - m^2 l^2 \right] \Phi_s = 0, \quad (34)$$

which is the same form as Eq.(10) when  $s = 0$ . On the other hand, in the far region, one has

$$\left[ \frac{d^2}{dx^2} + \frac{3}{x} \frac{d}{dx} - \frac{2s^2(x_+^2 - x_-^2)}{x^4(\cosh 2\delta + \cosh 2\gamma)} - \frac{m^2 l^2}{x^2} \right] \Phi_s^{\text{far}} = 0. \quad (35)$$

When  $s = 0$  or  $x \rightarrow \infty$ , this leads to Eq.(11). Hence, in the limit of  $x \rightarrow \infty (z \rightarrow \infty)$  the spin term does not play an important role. Also this is confirmed from the observation of Eq.(25) with  $\rho \rightarrow \infty$ .

In conclusion, we obtain the greybody factor for test fields  $\{\Phi_i\}$  on the  $AdS_3$ . This is based on both the non-normalizable mode of the test fields and  $AdS_3$ /CFT correspondence. Because the non-normalizable mode means the other type of boundary condition at infinity, we don't need to require the Dirichlet or Neumann condition. However, we wish to point out the ambiguity in determining normalization constant of the greybody factor in our  $AdS_3$ -calculation. It seems that this arises from the flux (13) which is calculated in the asymptotically  $AdS_3$ . The definition of flux is clear in the asymptotically flat space, but its definition seems to be unclear in the asymptotically  $AdS_3$ . For the test fields which couple to chiral operator, there exist some discrepancies in the greybody factor between the semiclassical and effective string calculations. We may resolve this by using the  $AdS_3$ /CFT correspondence. In the above we sketch the spin-dependent wave functions briefly, but we do not have a significant result until now. On the other hand, the greybody factor in (2) is valid for any  $h_L$  and  $h_R$ . Although (23) is derived for  $h_L = h_R$ , it takes the same form as (2). Hence we believe that our result (23) can be extended to  $h_L \neq h_R$  case to accommodate the test field with non-zero spin.

### ACKNOWLEDGEMENT

We would like to thank J.Y. Kim for usefull discussions. This work was supported in part by the Basic Science Research Institute Program, Minstry of Education, Project NOs. BSRI-98-2441 and BSRI-98-2413.

## REFERENCES

- [1] J. Maldacena, hep-th/9711200.
- [2] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, hep-th/9802109.
- [3] E. Witten, hep-th/9802150.
- [4] C. Callan, S. Gubser, I.G. Klebanov and A. Tseytlin, Nucl. Phys. **B489** (1997)65; M. Krasnitz and I.G. Klebanov, hep-th/9703216.
- [5] H.W. Lee, Y.S. Myung and J.Y. Kim, to appear in Phys. Rev. **D58**, hep-th/9708099.
- [6] I.R. Klebanov, A. Rajaraman and A. Tseytlin, Nucl. Phys. **B503**(1997) 157, hep-th/9704112.
- [7] J. Maldacena and A. Strominger, hep-th/9804085.
- [8] J. de Boer, hep-th/9806104.
- [9] S.S. Gubser, Phys. Rev. **D56**(1997) 7854; hep-th/9706100.
- [10] E. Teo, hep-th/9805014.
- [11] V. Balasubramanian, P. Kraus, and A. Lawrence, hep-th/9805171.
- [12] J.S. Chan and R.B. Mann, Phys. Rev. **D55**(1997)7546.
- [13] H.W. Lee, N.J. Kim, and Y.S. Myung, hep-th/9803227.
- [14] H.W. Lee, N.J. Kim, and Y.S. Myung, to appear in Phys. Rev. **D58**, hep-th/9803080.
- [15] H.W. Lee and Y.S. Myung, to appear in Phys. Rev. **D58**, hep-th/9804095.
- [16] H.W. Lee, N.J. Kim, and Y.S. Myung, hep-th/9805050.
- [17] M. Banados, C. Teitelboim and A. Zanelli, Phys. Rev. Lett. **69** (1992)1849; S. Carlip, Class. Quant. Grav. **12** (1995)2853.
- [18] D.B. Birmingham, I. Sachs and S. Sen, Phys. Lett. **B413** (1997)281.

- [19] J. Maldacena and A. Strominger, Phys. Rev. **D56**(1997)4975.
- [20] S. Carlip, hep-th/9806026.
- [21] K. Behrndt, I. Brunner, and I. Gaida, hep-th/9806195.